

# ABOUT THE LOGARITHM FUNCTION OVER THE MATRICES

GERALD BOURGEOIS

**ABSTRACT.** We prove the following results: let  $x, y$  be  $(n, n)$  complex matrices such that  $x, y, xy$  have no eigenvalue in  $] -\infty, 0]$  and  $\log(xy) = \log(x) + \log(y)$ . If  $n = 2$ , or if  $n \geq 3$  and  $x, y$  are simultaneously triangularizable, then  $x, y$  commute. In both cases we reduce the problem to a result in complex analysis.

## 1. INTRODUCTION

$\mathbb{Z}^*$  refers to the non-zero integers.

Let  $u$  be a complex number. Then  $Re(u), Im(u)$  refer to the real and imaginary parts of  $u$ ; if  $u \notin ] -\infty, 0]$  then  $arg(u) \in ] -\pi, \pi[$  refers to its principal argument.

**1.1. Basic facts about the logarithm.** Let  $x$  be a complex  $(n, n)$  matrix which hasn't any eigenvalue in  $] -\infty, 0]$ . Then  $\log(x)$ , the  $x$ -principal logarithm, is the  $(n, n)$  matrix  $a$  such that:

$e^a = x$  and the eigenvalues of  $a$  lie in the strip  $\{z \in \mathbb{C} : Im(z) \in ] -\pi, \pi[ \}$ .

$\log(x)$  always exists and is unique; moreover  $\log(x)$  may be written as a polynomial in  $x$ .

Now we consider two matrices  $x, y$  which have no eigenvalue in  $] -\infty, 0]$ :

- If  $x, y$  commute then  $x, y$  are simultaneously triangularizable and we may associate pairwise their eigenvalues  $(\lambda_j), (\mu_j)$ ; if moreover  $\forall j, |arg(\lambda_j) + arg(\mu_j)| < \pi$ , then  $\log(xy) = \log(x) + \log(y)$ .
- Conversely if  $xy$  has no eigenvalue in  $] -\infty, 0]$  and  $\log(xy) = \log(x) + \log(y)$  then do  $x, y$  commute? We will prove that it's true for  $n = 2$  (theorem 1) or, for all  $n$ , if  $x, y$  are simultaneously triangularizable (theorem 2). But if  $n > 2$ , then we don't know the answer in the general case.

**1.2. Lemma 1.** Let  $x, y$  be two complex  $(n, n)$  matrices such that  $x, y$  haven't any eigenvalue in  $] -\infty, 0]$  and  $\log(x)\log(y) = \log(y)\log(x)$ .

Then  $x, y$  commute.

**Proof.** The principal logarithm over  $\mathbb{C} \setminus ] -\infty, 0]$  is one to one; thus, using Hermite's interpolation formula,  $x$  or  $y$  may be written as a polynomial in  $\log(x)$  or  $\log(y)$ .  $\square$

---

2000 *Mathematics Subject Classification.* Primary 39B42.

*Key words and phrases.* Linear algebra, matrix, logarithm.

## 2. DIMENSION 2

**2.1. Principle of the proof.** The proof is based on the two next propositions. The first one is a corollary of a Morinaga and Nono's result ([1, p. 356]); the second is a technical result using complex analysis.

**Proposition 1.** Let  $\mathcal{U} = \{u \in \mathbb{C}^* : e^u = 1 + u\}$ .

Let  $a, b$  be two  $(2, 2)$  complex matrices such that  $e^{a+b} = e^a e^b$  and  $ab \neq ba$ ; let  $\text{spectrum}(a) = \{\lambda_1, \lambda_2\}$ ,  $\text{spectrum}(b) = \{\mu_1, \mu_2\}$ .

Then one of the three following *item* is fulfilled:

- (1)  $\lambda_1 - \lambda_2 \in 2i\pi\mathbb{Z}^*$  and  $\mu_1 - \mu_2 \in 2i\pi\mathbb{Z}^*$ .
- (2) One of the following complex numbers  $\pm(\lambda_1 - \lambda_2)$ ,  $\pm(\mu_1 - \mu_2)$  is in  $\mathcal{U}$ .
- (3)  $a$  and  $b$  are simultaneously similar to  $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda + u \end{pmatrix}$  and  $\begin{pmatrix} \mu + v & 1 \\ 0 & \mu \end{pmatrix}$  with  $\lambda, \mu \in \mathbb{C}$ ,  $u, v \in \mathbb{C}^*$ ,  $u \neq v$  and  $\frac{e^u - 1}{u} = \frac{e^v - 1}{v} \neq 0$ .

**Proposition 2.** Let  $u, v$  be two distinct, non zero complex numbers such that  $\frac{e^u - 1}{u} = \frac{e^v - 1}{v} \neq 0$ ,  $|Im(u)| < 2\pi$ ,  $|Im(v)| < 2\pi$ . Then necessarily  $|Im(u) - Im(v)| \geq 2\pi$ .

**Proof.** Assume that we can choose these  $u, v$  such that  $|Im(u) - Im(v)| < 2\pi$ . Let  $\lambda = \frac{e^u - 1}{u}$  and let  $f$  be the holomorphic function:  $f(z) = e^z - \lambda z - 1$ .

Now we show that there exists  $a \in ]0, 2\pi[$  such that  $Im(u), Im(v)$  are in  $] -a, 2\pi - a[$  and  $f$  hasn't any zero with imaginary part  $-a$  or  $2\pi - a$ .

If it's false then  $f$  admits an infinity of zeros in the strip  $\{z : Im(z) \in ] -2\pi, 2\pi[ \}$ :

Case 1: we can extract a sequence of zeros  $z_k$  such that  $Re(z_k) \rightarrow -\infty$ ; then  $f(z_k) \sim -\lambda z_k$ , a contradiction.

Case 2: we can extract a sequence of zeros  $z_k$  such that  $Re(z_k) \rightarrow +\infty$ ; then  $f(z_k) \sim e^{z_k}$ , a contradiction.

Let  $r$  be a big positive real such that  $Re(u), Re(v)$  are in  $] -r, r[$  and  $\Delta$  be the rectangle  $\{z : -r \leq Re(z) \leq r, -a \leq Im(z) \leq 2\pi - a\}$ . The oriented edge  $\partial\Delta$  consists of four parts:  $h_1 = \{x + i(2\pi - a) : x \text{ from } r \text{ to } -r\}$ ,  $v_1 = \{-r + iy : y \text{ from } 2\pi - a \text{ to } -a\}$ ,  $h_2 = \{x - ia : x \text{ from } -r \text{ to } r\}$ ,  $v_2 = \{r + iy : y \text{ from } -a \text{ to } 2\pi - a\}$ .  $f$

admits in  $\Delta$  at least three zeros:  $0, u, v$ . Thus  $\frac{1}{2i\pi} \int_{\partial\Delta} \frac{f'(z)}{f(z)} dz = I(f(\partial\Delta), 0) \geq 3$  where  $I$  refers to the index function.

$f(z + 2i\pi) - f(z) = -\lambda 2i\pi$ ; then  $\{f(h_1), f(h_2)\}$  is in tubular form; moreover  $f(h_1)$  and  $f(h_2)$  are isometric to a parametric curve in the form  $\{(e^t - \sigma t, \tau t) : t \in [-r, r]\}$  where  $\sigma, \tau$  are real; we can choose  $a$  such that  $\sigma, \tau \in \mathbb{R}^*$ ; thus for  $j \in \{1, 2\}$

$$\left| \frac{1}{2i\pi} \int_{h_j} \frac{f'(z)}{f(z)} dz \right| \leq 1 - \rho \text{ with } \rho = \frac{1}{2\pi} \arctan\left(\left|\frac{\tau}{\sigma}\right|\right).$$

We can choose  $r, a$  such that  $f'(z) \neq 0$  on  $\partial\Delta$ ; thus  $f(h_1), f(h_2)$  intersect perpendicularly  $f(v_1)$  and  $f(v_2)$ . If  $z \in v_1$  and  $r \rightarrow +\infty$  then  $f(z) = -\lambda z - 1 + O(e^{-r})$ ;  $f(v_1)$  is close to a segment of fixed direction and length. If  $z \in v_2$  and  $r \rightarrow +\infty$  then  $f(z) = e^z + O(r)$ ;  $f(v_2)$  is close to an anticlockwise circle of radius  $e^r$  containing 0.

Therefore  $\frac{1}{2i\pi}(\int_{h_1} \frac{f'(z)}{f(z)}dz + \int_{h_2} \frac{f'(z)}{f(z)}dz) \leq 2 - 2\rho$ ,  $\frac{1}{2i\pi}(\int_{v_1} \frac{f'(z)}{f(z)}dz) \approx 0$ ,  $\frac{1}{2i\pi}(\int_{v_2} \frac{f'(z)}{f(z)}dz) \approx 1$ ; what is contradictory with  $I(f(\partial\Delta), 0) \geq 3$ .  $\square$

**2.2. Theorem 1.** Let  $x, y$  be two  $(2, 2)$  complex matrices such that  $x, y, xy$  haven't any eigenvalue in  $]-\infty, 0]$  and  $\log(xy) = \log(x) + \log(y)$ . Then  $x, y$  commute.

**Proof.** We assume that  $xy \neq yx$ .  $e^{\log(x)}e^{\log(y)} = e^{\log(x)+\log(y)}$ ; using lemma 1,  $\log(x)\log(y) \neq \log(y)\log(x)$ ; thus we may use Proposition 1; it's wellknown that  $u \in \mathcal{U}$  implies that  $|Im(u)| > 2\pi$ ; then, according to the logarithm definition,  $a = \log(x)$  and  $b = \log(y)$  satisfy *item* (3). Moreover the conditions  $|Im(u)| < 2\pi, |Im(v)| < 2\pi, |Im(u) - Im(v)| < 2\pi$  are necessarily fulfilled. Proposition 2 proves that these conditions can't be all satisfied.  $\square$

### 3. DIMENSION $n$

$I$  refers to the identity matrix of dimension  $n - 1$ . Let  $\phi$  be the holomorphic function:  $\phi : z \rightarrow \frac{e^z - 1}{z}, \phi(0) = 1$ .

**Remark 1.** We have shown in part 2 that if  $u, v$  are complex numbers such that  $|Im(u)| < 2\pi, |Im(v)| < 2\pi, |Im(u - v)| < 2\pi$  and  $\phi(u) = \phi(v)$ , then  $u = v$ . We'll use the following to prove our second main result.

**3.1. Proposition 3.** Let  $a = \begin{pmatrix} a_0 & u \\ 0 & \alpha \end{pmatrix}, b = \begin{pmatrix} b_0 & v \\ 0 & \beta \end{pmatrix}$  be two complex  $(n, n)$  matrices where  $\alpha, \beta$  are complex numbers and  $a_0, b_0$  are  $(n - 1, n - 1)$  complex matrices which commute; let  $spectrum(a_0 - \alpha I) = (\alpha_i)_{i \leq n-1}, spectrum(b_0 - \beta I) = (\beta_i)_{i \leq n-1}$ . If  $e^{a+b} = e^a e^b$  and  $ab \neq ba$  then one of the following *item* must be satisfied:

- (4)  $\exists i : \beta_i \neq 0$  and  $\phi(\alpha_i + \beta_i) = \phi(\alpha_i)$ .
- (5)  $\exists i : \alpha_i \neq 0, \beta_i = 0$  and  $\phi(-\alpha_i) = 1$ .

**Proof.** We may assume that  $a_0, b_0$  are upper triangular. Let  $a_1 = a_0 - \alpha I, b_1 = b_0 - \beta I, w = a_1 v - b_1 u = [w_1, \dots, w_{n-1}]^T$ . Thus  $ab \neq ba$  iff  $w \neq 0$ .

Then  $e^a = e^\alpha \begin{pmatrix} e^{a_1} & \phi(a_1)u \\ 0 & 1 \end{pmatrix}, e^b = e^\beta \begin{pmatrix} e^{b_1} & \phi(b_1)v \\ 0 & 1 \end{pmatrix},$

$e^{a+b} = e^{\alpha+\beta} \begin{pmatrix} e^{a_1+b_1} & \phi(a_1+b_1)(u+v) \\ 0 & 1 \end{pmatrix}$ ; therefore  $e^{a+b} = e^a e^b$  iff

- (6)  $(\phi(a_1 + b_1) - \phi(a_1))u = (e^{a_1}\phi(b_1) - \phi(a_1 + b_1))v$ .  
 $(e^{a_1}\phi(b_1) - \phi(a_1 + b_1))b_1 = (\phi(a_1 + b_1) - \phi(a_1))a_1$  and (6) imply that  $(e^{a_1}\phi(b_1) - \phi(a_1 + b_1))b_1 v = (\phi(a_1 + b_1) - \phi(a_1))a_1 v = (\phi(a_1 + b_1) - \phi(a_1))b_1 u$ ; thus
- (7)  $(\phi(a_1 + b_1) - \phi(a_1))w = 0$ .

We have also  $e^b = e^{-a} e^{a+b}$ ; then we can prove by the same method that

- (8)  $(\phi(b_1) - \phi(-a_1))w = 0$ .

There exists  $k$  such that  $w_k \neq 0$  and if  $j > k$  then  $w_j = 0$ . Therefore (7),(8) imply that  $\phi(\alpha_k + \beta_k) = \phi(\alpha_k)$  and  $\phi(\beta_k) = \phi(-\alpha_k)$ ; we are done except if  $\alpha_k = \beta_k = 0$ .

Now we assume that  $\alpha_k = \beta_k = 0$ .  $\phi(a_1 + b_1) - \phi(a_1) = \frac{1}{2}b_1(I + P(a_1, b_1)), e^{a_1}\phi(b_1) - \phi(a_1 + b_1) = \frac{1}{2}a_1(I + P(a_1, b_1))$  where  $P$  is an analytic function, defined on  $\mathbb{C}^2$ ,

which satisfies  $P(0, 0) = 0$ . (6) can be rewritten as  $(I + P(a_1, b_1))w = 0$ . Therefore  $(1 + P(0, 0))w_k = 0$ , a contradiction.  $\square$

**3.2. Theorem 2.** Let  $x, y$  be  $(n, n)$  complex matrices such that  $x, y, xy$  haven't any eigenvalue in  $] - \infty, 0]$  and  $\log(xy) = \log(x) + \log(y)$ . If moreover  $x, y$  are simultaneously triangularizable then  $xy = yx$ .

**Proof.** We assume that  $x, y$  are upper-triangular and  $xy \neq yx$ ; we prove inductively the result for  $n \geq 2$ .  $x = \begin{pmatrix} x_0 & ? \\ 0 & \lambda \end{pmatrix}$ ,  $y = \begin{pmatrix} y_0 & ? \\ 0 & \mu \end{pmatrix}$  where  $x_0, y_0$  are  $(n-1, n-1)$  upper triangular matrices which haven't any eigenvalue in  $] - \infty, 0]$  and  $\lambda, \mu \in \mathbb{C} \setminus ] - \infty, 0]$ . The matrices  $a = \log(x), b = \log(y)$  are polynomials in  $x$  or  $y$ , thus they are upper-triangular in form  $a = \begin{pmatrix} \log(x_0) & ? \\ 0 & \log(\lambda) \end{pmatrix}$ ,  $b = \begin{pmatrix} \log(y_0) & ? \\ 0 & \log(\mu) \end{pmatrix}$ . Thus  $\log(x_0 y_0) = \log(x_0) + \log(y_0)$ ; according to the recurrence hypothesis  $x_0 y_0 = y_0 x_0$  and then  $\log(x_0) \log(y_0) = \log(y_0) \log(x_0)$ . Moreover  $e^{a+b} = e^a e^b$  and, from lemma 1,  $ab \neq ba$ .

Now we use Proposition 3 with  $\alpha = \log(\lambda), \beta = \log(\mu), a_0 = \log(x_0), b_0 = \log(y_0)$ . Here  $\alpha_i, \beta_i, \alpha_i + \beta_i$  have imaginary parts in  $] - 2\pi, 2\pi[$  and according to Remark 1, item (4),(5) can't be satisfied.  $\square$

We conclude with an easy result.

**3.3. Proposition 4.** Let  $x, y$  be two positive definite hermitian  $(n, n)$  matrices so that  $\log(xy) = \log(x) + \log(y)$ . Then  $xy = yx$ .

**Proof.**  $\log(xy)$  exists because  $\text{spectrum}(xy) \subset ]0, \infty[$ ;  $a = \log(x), b = \log(y)$  are hermitian matrices such that  $e^{a+b} = e^a e^b$ . Moreover  $e^{a+b} = (e^{a+b})^* = e^b e^a$  and  $e^a e^b = e^b e^a$  or  $xy = yx$ .  $\square$

**Remark.** It's wellknown that if  $a, b$  are bounded self adjoint operators on a complex Hilbert space, then  $e^{a+b} = e^a e^b$  implies that  $ab = ba$ . ( cf. [2, Corollary 1]).

#### 4. CONCLUSION

When  $n = 2$ , we know how to characterize the complex  $(n, n)$  matrices  $a, b$  such that  $ab \neq ba$  and  $e^{a+b} = e^a e^b$ ; it allowed us to bring back our problem to a result of complex analysis. Unfortunately, if  $n \geq 3$ , the classification of such matrices is unknown. For this reason we can't prove, in this last case, the hoped result without supplementary assumption.

**Acknowledgement.** The author would like to thank F. Nazarov for his participation in the proof of proposition 2.

#### REFERENCES

- [1] K. Morinaga and T.Nono. On the non-commutative solutions of the exponential equation  $e^x e^y = e^{x+y}$ . J. Sci. Hiroshima univ. (A)17, (1954), 345-358.

- [2] C. Schmoeger. Remarks on commuting exponentials in Banach algebras. Proceedings of the American Mathematical Society. Volume 127, Number 5, (1999), pages 1337,1338.

DEPARTEMENT DE MATHÉMATIQUES, FACULTE DE LUMINY, 163 AVENUE DE LUMINY, CASE 901,  
13288 MARSEILLE CEDEX 09, FRANCE.

*E-mail address:* `bourgeoi@luminath.univ-mrs.fr`